

This article was downloaded by: [Marco Fontana]

On: 26 September 2011, At: 06:11

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lagb20>

An Overring-Theoretic Approach to Polynomial Extensions of Star and Semistar Operations

Gyu Whan Chang^a & Marco Fontana^b

^a Department of Mathematics, University of Incheon, Incheon, Korea

^b Dipartimento di Matematica, Università degli Studi "Roma Tre", Rome, Italy

Available online: 21 Jun 2011

To cite this article: Gyu Whan Chang & Marco Fontana (2011): An Overring-Theoretic Approach to Polynomial Extensions of Star and Semistar Operations, Communications in Algebra, 39:6, 1956-1978

To link to this article: <http://dx.doi.org/10.1080/00927872.2010.480959>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan, sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN OVERRING-THEORETIC APPROACH TO POLYNOMIAL EXTENSIONS OF STAR AND SEMISTAR OPERATIONS

Gyu Whan Chang¹ and Marco Fontana²

¹Department of Mathematics, University of Incheon, Incheon, Korea

²Dipartimento di Matematica, Università degli Studi “Roma Tre”, Rome, Italy

Call a semistar operation \circledast on the polynomial domain $D[X]$ an extension (respectively, a strict extension) of a semistar operation \star defined on an integral domain D , with quotient field K , if $E^\star = (E[X])^\circledast \cap K$ (respectively, $E^\star[X] = (E[X])^\circledast$) for all nonzero D -submodules E of K . In this article, we study the general properties of the above defined extensions and link our work with earlier efforts, centered on the stable semistar operation case, at defining semistar operations on $D[X]$ that are “canonical” extensions (or, “canonical” strict extensions) of semistar operations on D .

Key Words: Kronecker function ring; Localizing system of ideals; Nagata ring; Polynomial domain; Star and semistar operation; v -, t -, w -, b -operation.

2000 Mathematics Subject Classification: 13B25; 13A15; 13G05; 13B22.

1. BACKGROUND RESULTS

Let D be an integral domain with quotient field K . Let $\overline{F}(D)$ denote the set of all nonzero D -submodules of K , and let $F(D)$ be the set of all nonzero fractional ideals of D , i.e., $E \in F(D)$ if $E \in \overline{F}(D)$, and there exists a nonzero $d \in D$ with $dE \subseteq D$. Let $f(D)$ be the set of all nonzero finitely generated D -submodules of K . Then, obviously $f(D) \subseteq F(D) \subseteq \overline{F}(D)$.

Following Okabe and Matsuda [28], a *semistar operation* on D is a map $\star : \overline{F}(D) \rightarrow \overline{F}(D)$, $E \mapsto E^\star$, such that, for all $x \in K$, $x \neq 0$, and for all $E, F \in \overline{F}(D)$, the following properties hold:

- (\star_1) $(xE)^\star = xE^\star$;
- (\star_2) $E \subseteq F$ implies $E^\star \subseteq F^\star$;
- (\star_3) $E \subseteq E^\star$ and $E^{\star\star} := (E^\star)^\star = E^\star$.

The semistar operation defined by $E^\star = K$ for all $E \in \overline{F}(D)$ is called the *trivial semistar operation on D* , and it is denoted by e_D (or, simply, by e). A *(semi)star operation* is a semistar operation that, restricted to $F(D)$, is a star operation (in the sense of [12, Section 32]). It is easy to see that a semistar operation \star on D is a (semi)star operation if and only if $D^\star = D$.

Received November 30, 2009; Revised February 12, 2010. Communicated by T. Albu.

Address correspondence to Marco Fontana, Dipartimento di Matematica, Università degli Studi “Roma Tre”, Rome 00146, Italy; E-mail: fontana@mat.uniroma3.it

If \star is a semistar operation on D , then we can consider a map $\star_f : \overline{F}(D) \rightarrow \overline{F}(D)$ defined, for each $E \in \overline{F}(D)$, as follows:

$$E^{\star_f} := \bigcup \{F^{\star} \mid F \in f(D) \text{ and } F \subseteq E\}.$$

It is easy to see that \star_f is a semistar operation on D , called *the semistar operation of finite type associated to \star* . Note that, for each $F \in f(D)$, $F^{\star} = F^{\star_f}$. A semistar operation \star is called a *semistar operation of finite type* if $\star = \star_f$. It is easy to see that $(\star_f)_f = \star_f$ (that is, \star_f is of finite type).

If \star_1 and \star_2 are two semistar operations on D , we say that $\star_1 \leq \star_2$ if $E^{\star_1} \subseteq E^{\star_2}$, for each $E \in \overline{F}(D)$. This is equivalent to say that $(E^{\star_1})^{\star_2} = E^{\star_2} = (E^{\star_2})^{\star_1}$, for each $E \in \overline{F}(D)$. Obviously, for each semistar operation \star , we have $\star_f \leq \star$. Let d_D (or, simply, d) be the *identity (semi)star operation on D* ; clearly, $d \leq \star$, for all semistar operation \star on D .

We say that a nonzero ideal I of D is a *quasi- \star -ideal* if $I^{\star} \cap D = I$, a *quasi- \star -prime ideal* if it is a prime quasi- \star -ideal, and a *quasi- \star -maximal ideal* if it is maximal in the set of all proper quasi- \star -ideals. A quasi- \star -maximal ideal is a prime ideal. It is possible to prove that each proper quasi- \star_f -ideal is contained in a quasi- \star_f -maximal ideal. More details can be found in [10, p. 4781]. We will denote by $\text{QMax}^{\star}(D)$ (respectively, $\text{QSpec}^{\star}(D)$) the set of the quasi- \star -maximal ideals (respectively, quasi- \star -prime ideals) of D . When \star is a (semi)star operation, the notion of quasi- \star -ideal coincides with the “classical” notion of \star -ideal (i.e., a nonzero ideal I such that $I^{\star} = I$).

If Δ is a set of prime ideals of an integral domain D , then the semistar operation \star_{Δ} defined on D

$$E^{\star_{\Delta}} := \bigcap \{ED_P \mid P \in \Delta\}, \quad \text{for each } E \in \overline{F}(D),$$

is called *the spectral semistar operation on D associated to Δ* . A semistar operation \star on an integral domain D is called a *spectral semistar operation* if there exists a subset Δ of the prime spectrum of D , $\text{Spec}(D)$, such that $\star = \star_{\Delta}$.

When $\Delta := \text{QMax}^{\star_f}(D)$, we set $\tilde{\star} := \star_{\Delta}$, i.e.,

$$E^{\tilde{\star}} := \bigcap \{ED_P \mid P \in \text{QMax}^{\star_f}(D)\}, \quad \text{for each } E \in \overline{F}(D).$$

A semistar operation \star is *stable* if $(E \cap F)^{\star} = E^{\star} \cap F^{\star}$, for each $E, F \in \overline{F}(D)$. Spectral semistar operations are stable [6, Lemma 4.1(3)]. In particular, $\tilde{\star}$ is a semistar operation stable and of finite type; and, conversely, if a semistar operation \star is stable and of finite type then $\star = \tilde{\star}$ [6, Corollary 3.9(2)].

Let T be an overring of an integral domain D , let $\iota : D^{\times} \rightarrow T$ be the canonical embedding, and let \star be a semistar operation on D . We denote by \star_{ι} the semistar operation on T defined by $E^{\star_{\iota}} := E^{\star}$, for each $E \in \overline{F}(T) (\subseteq \overline{F}(D))$.

Conversely, let \ast be a semistar operation on T , and let \ast' be the semistar operation on D defined by $E^{\ast'} := (ET)^{\ast}$, for each $E \in \overline{F}(D)$.

It is not difficult to see that $(\ast')_f = (\ast_f)'$ and if \star is a semistar operation of finite type (respectively, a stable semistar operation) on D , then \star_{ι} is a semistar operation of finite type (respectively, a stable semistar operation) on T (cf. for instance [9, Proposition 2.8] and [29, Propositions 2.11 and 2.13]).

Clearly, if $\star = d_D$, then $(d_D)_t = d_T$. In case $\star = d_T$, the semistar operation of finite type $(d_T)'$ (defined by $E \mapsto ET$ for all $E \in \overline{F}(D)$) is denoted also by $\star_{\{T\}}$, and it is stable if and only if T is a flat overring of D [34, Proposition 1.7] and [25, Theorem 7.4(1)].

By v_D (or, simply, by v), we denote the v -(semi)star operation defined as usual by $E^v := (D : (D : E))$, for each $E \in \overline{F}(D)$. By t_D (or, simply, by t) we denote $(v_D)_f$ the t -(semi)star operation on D and by w_D (or just by w) the stable semistar operation of finite type associated to v_D (or, equivalently, to t_D), considered by Wang and McCasland in [35] (cf. also [13]); i.e., $w_D := \tilde{v}_D = \tilde{t}_D$. Clearly, $w_D \leq t_D \leq v_D$. Moreover, it is easy to see that for each (semi)star operation \star on D , we have $\star \leq v_D$ and $\star_f \leq t_D$ (cf. also [12, Theorem 34.1(4)]).

We recall from [7, Chapter V] (see also [31, Chapter 4]) that a *localizing system of ideals of D* is a family \mathcal{F} of ideals of D such that:

- (LS1) If $I \in \mathcal{F}$ and J is an ideal of D such that $I \subseteq J$, then $J \in \mathcal{F}$;
- (LS2) If $I \in \mathcal{F}$ and J is an ideal of D such that $(J :_D iD) \in \mathcal{F}$, for each $i \in I$, then $J \in \mathcal{F}$.

A localizing system \mathcal{F} is *finitely generated* if, for each $I \in \mathcal{F}$, there exists a finitely generated ideal $J \in \mathcal{F}$ such that $J \subseteq I$.

The relation between stable semistar operations and localizing systems has been investigated by Fontana and Huckaba in [6] and by Halter-Koch in the context of module systems [17]. In the following proposition, we summarize some of the results that we need (see [6, Proposition 2.8, Proposition 3.2, Proposition 2.4, Corollary 2.11, Theorem 2.10(B)]).

Proposition 1. *Let D be an integral domain.*

- (1) If \star is a semistar operation on D , then $\mathcal{F}^\star := \{I \text{ ideal of } D \mid I^\star = D^\star\}$ is a localizing system (called the localizing system associated to \star).
- (2) If \star is a semistar operation of finite type, then \mathcal{F}^\star is a finitely generated localizing system.
- (3) Let $\star_{\mathcal{F}}$ or, simply, $\overline{\star}$ be the semistar operation associated to a given localizing system \mathcal{F} of D and defined by $E \mapsto E^\star := \bigcup \{(E : J) \mid J \in \mathcal{F}\}$, for each $E \in \overline{F}(D)$. Then $\star_{\mathcal{F}}$ (called the semistar operation associated to the localizing system \mathcal{F}) is a stable semistar operation on D .
- (4) $\overline{\star} \leq \star$ and $\mathcal{F}^\star = \mathcal{F}^{\overline{\star}}$.
- (5) $\overline{\star} = \star$ if and only if \star is stable.
- (6) If \mathcal{F} is a finitely generated localizing system, then $\star_{\mathcal{F}}$ is a finite type (stable) semistar operation.
- (7) $\mathcal{F}^{\star_f} = (\mathcal{F}^\star)_f := \{I \in \mathcal{F}^\star \mid I \supseteq J, \text{ for some finitely generated ideal } J \in \mathcal{F}^\star\}$ and $\tilde{\star} = \overline{\star}_f$, i.e., $\tilde{\star}$ is the stable semistar operation of finite type associated to the localizing system \mathcal{F}^{\star_f} . In particular, for each $E \in \overline{F}(D)$, we have:

$$E^{\tilde{\star}} = \bigcup \{(E : J) \mid J \in \mathcal{F}(D), J \subseteq D, \text{ and } J^\star = D^\star\}.$$

- (8) If \mathcal{F}' and \mathcal{F}'' are two localizing systems of D , then $\mathcal{F}' \subseteq \mathcal{F}''$ if and only if $\star_{\mathcal{F}'} \leq \star_{\mathcal{F}''}$.

If I is a nonzero fractional ideal of D , we say that I is \star -invertible if $(II^{-1})^\star = D^\star$. From the definitions and from the fact that $\text{QMax}^{\star_f}(D) = \text{QMax}^\star(D)$ [10, Corollary 3.5(2)] it follows easily that a nonzero fractional ideal I is $\tilde{\star}$ -invertible if and only if I is \star_f -invertible. An integral domain D is called a *Prüfer \star -multiplication domain* (for short, *P \star MD*) if each $I \in \mathbf{f}(D)$ is \star_f -invertible. It is easy to see that the notions of PvMD, PtMD and PwMD coincide. Obviously, a PdMD is a Prüfer domain, and conversely [12, Theorem 22.1].

If R is a ring (not necessarily an integral domain) and X an indeterminate over R , then the ring $R(X) := \{f/g \mid f, g \in R[X] \text{ and } c(g) = R\}$ (where $c(g)$ is the content of the polynomial g) is called the *Nagata ring* of R [12, Proposition 33.1].

In case of an integral domain equipped with a semistar operation, we have a general “semistar version” of the Nagata ring. The following result was proved in [10, Propositions 3.1 and 3.4] (cf. also [23, Proposition 2.1]).

Proposition 2. *Let \star be a nontrivial semistar operation on an integral domain D . Set $\mathcal{N}^\star := \mathcal{N}_D^\star := \{h \in D[X] \mid h \neq 0 \text{ and } c(h)^\star = D^\star\}$ and*

$$Na(D, \star) := D[X]_{\mathcal{N}^\star}.$$

Then,

- (1) \mathcal{N}^\star is a saturated multiplicative subset of $D[X]$ and $\mathcal{N}^\star = \mathcal{N}^{\star_f} = D[X] \setminus \bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\}$.
- (2) $\text{Max}(Na(D, \star)) = \{Q[X]_{\mathcal{N}^\star} \mid Q \in \text{QMax}^{\star_f}(D)\}$ and $\text{QMax}^{\star_f}(D)$ coincides with the canonical image in $\text{Spec}(D)$ of $\text{Max}(Na(D, \star))$.
- (3) $Na(D, \star) = \bigcap \{D_Q(X) \mid Q \in \text{QMax}^{\star_f}(D)\}$.
- (4) For each $E \in \overline{\mathbf{F}}(D)$, $E^\star = ENa(D, \star) \cap K$.

Let \star be a semistar operation on D . If F is in $\mathbf{f}(D)$, we say that F is \star -eab (respectively, \star -ab) if $(FG)^\star \subseteq (FH)^\star$ implies that $G^\star \subseteq H^\star$, with $G, H \in \mathbf{f}(D)$, (respectively, with $G, H \in \overline{\mathbf{F}}(D)$).

An operation \star is said to be eab (respectively, ab) if each $F \in \mathbf{f}(D)$ is \star -eab (respectively, \star -ab). An ab operation is obviously an eab operation. We note that if \star is an eab semistar operation, then \star_f is also an eab semistar operation, since they agree on all finitely generated ideals. Let \star be a semistar operation of finite type. Then \star is an eab semistar operation if and only if \star is an ab semistar operation. In this situation, we say that \star is an *(e)ab semistar operation*. In particular, from the previous result, it follows that the notions of \star -eab semistar operation and \star_f -(e)ab semistar operation coincide [11, Lemma 3 and Proposition 4].

Given an arbitrary semistar operation \star on an integral domain D , it is possible to associate to \star , an eab semistar operation of finite type \star_a of D , called the ab semistar operation associated to \star , defined as

$$F^{\star_a} := \bigcup \{((FH)^\star : H) \mid H \in \mathbf{f}(D)\}, \quad \text{for each } F \in \mathbf{f}(D),$$

and, in general,

$$E^{\star_a} := \bigcup \{F^{\star_a} \mid F \subseteq E, F \in \mathbf{f}(D)\}, \quad \text{for each } E \in \overline{\mathbf{F}}(D),$$

[9, Definition 4.4]. Note that if \star is an (e) **ab** semistar operation of finite type, then $\star = \star_a$, and conversely [9, Proposition 4.5]. More information on the operation \star_a , introduced for ideal systems in [22, p. 41] (see also Lorenzen's original article [24]) can be found in [8, 15, 16, 27, 28].

Let \star be a semistar operation on D , and let V be a valuation overring of D . We say that V is a \star -valuation overring of D if, for each $F \in \mathbf{f}(D)$, $F^\star \subseteq FV$ (or equivalently, $\star_f \leq \star_{\{V\}}$). Note that a valuation overring V of D is a \star -valuation overring of D if and only if $V^{\star_f} = V$. More details of semistar valuation overrings can be found in [8, 9] (cf. also [15, 18, 22]).

Proposition 3. ([9, Proposition 3.3, Theorem 3.11, Theorem 5.1, Corollary 5.2, Corollary 5.3], [8, Theorem 3.5]). *Let \star be any semistar operation defined on an integral domain D with quotient field K and let \star_a be the **ab** semistar operation associated to \star . Set*

$$Kr(D, \star) := \{f/g \mid f, g \in D[X] \setminus \{0\} \text{ and there exists } h \in D[X] \setminus \{0\} \\ \text{such that } (c(f)c(h))^\star \subseteq (c(g)c(h))^\star\} \cup \{0\}.$$

Then, we have:

- (1) $Kr(D, \star)$ is a Bézout domain with quotient field $K(X)$, called the Kronecker function ring of D with respect to the semistar operation \star ;
- (2) $Na(D, \star) \subseteq Kr(D, \star)$;
- (3) $Kr(D, \star) = Kr(D, \star_a)$;
- (4) $E^{\star_a} = EKr(D, \star) \cap K$, for each $E \in \overline{F}(D)$;
- (5) $Kr(D, \star) = \bigcap \{V(X) \mid V \text{ is a } \star\text{-valuation overring of } D\}$.
- (6) If $F := (a_0, a_1, \dots, a_n) \in \mathbf{f}(D)$ and $f(X) := a_0 + a_1X + \dots + a_nX^n \in K[X]$, then

$$FKr(D, \star) = f(X)Kr(D, \star) = c(f)Kr(D, \star).$$

When $\star = d$, the d -valuation overrings of D are just the valuation overrings of D . In this case, we set

$$Kr(D) := Kr(D, d) = \bigcap \{V(X) \mid V \text{ is a valuation overring of } D\}.$$

Moreover, if we denote by b_D (or, simply, by b) the **ab** semistar operation of finite type $(d_D)_a$, then for each $E \in \overline{F}(D)$,

$$E^b = EKr(D) \cap K = \bigcap \{EV \mid V \text{ is a valuation overring of } D\}.$$

Remark 4. Recall that a Prüfer domain D can be characterized by the fact that each $F \in \mathbf{f}(D)$ is invertible. Since an invertible ideal is always a v -ideal (and, in particular, a t -ideal), then the following are equivalent [12, Theorem 24.7 and Theorem 34.1(4)]:

- (i) D is a Prüfer domain;
- (ii) D is integrally closed and $d = b$;
- (iii) D is integrally closed and $d = t$.

2. RESULTS

Let D be an integral domain with quotient field K , and let X be an indeterminate over K . We start with some basic facts. Note that some of the statements contained in the following result were also proved in [26, Proposition 2.1] and, in the star operation setting, in [20, Propositions 2.1 and 2.2].

Lemma 5. *Given a semistar operation \otimes on $D[X]$, for each $E \in \overline{F}(D)$ set*

$$E^{\otimes_0} := (E[X])^{\otimes} \cap K.$$

Then:

- (1) \otimes_0 is a semistar operation on D called the semistar operation canonically induced by \otimes on D . In particular, if \otimes is a (semi)star operation on $D[X]$, then \otimes_0 is a (semi)star operation on D ;
- (2) $(E^{\otimes_0}[X])^{\otimes} = (E[X])^{\otimes}$ for all $E \in \overline{F}(D)$;
- (3) $(\otimes_f)_0 = (\otimes_0)_f$;
- (4) If \otimes is a semistar operation of finite type (respectively, stable), then \otimes_0 is a semistar operation of finite type (respectively, stable);
- (5) If \otimes' and \otimes'' are two semistar operations on $D[X]$ and $\otimes' \leq \otimes''$, then $\otimes'_0 \leq \otimes''_0$;
- (6) $(\widetilde{\otimes})_0 = \widetilde{\otimes_0}$;
- (7) $(d_{D[X]})_0 = d_D$, $(w_{D[X]})_0 = w_D$, $(t_{D[X]})_0 = t_D$, $(v_{D[X]})_0 = v_D$, and $(b_{D[X]})_0 = b_D$.

Proof. (1) Set $\star := \otimes_0$. It is easy to see that, if $E \in \overline{F}(D)$, then $E \subseteq E^\star$, and if $E_1, E_2 \in \overline{F}(D)$ with $E_1 \subseteq E_2$, then $E_1^\star \subseteq E_2^\star$. Moreover,

$$\begin{aligned} (E^\star)^\star &= (((E[X])^{\otimes} \cap K)[X])^{\otimes} \cap K \\ &\subseteq ((E[X])^{\otimes}[X] \cap K[X])^{\otimes} \cap K \\ &= ((E[X])^{\otimes} \cap K[X])^{\otimes} \cap K \\ &\subseteq ((E[X])^{\otimes})^{\otimes} \cap K = (E[X])^{\otimes} \cap K = E^\star. \end{aligned}$$

Thus $(E^\star)^\star = E^\star$. Moreover, for each nonzero $z \in K$, we have

$$\begin{aligned} zE^\star &= z((E[X])^{\otimes} \cap K) \\ &= (z(E[X])^{\otimes} \cap zK) = (z(E[X])^{\otimes} \cap K) \\ &= (zE[X])^{\otimes} \cap K = (zE)^\star. \end{aligned}$$

In particular, if $(D[X])^{\otimes} = D[X]$, then $D^\star = (D[X])^{\otimes} \cap K = D[X] \cap K = D$.

(2) Note that $E[X] \subseteq E^{\otimes_0}[X] = ((E[X])^{\otimes} \cap K)[X] \subseteq (E[X])^{\otimes} \cap K[X] \subseteq (E[X])^{\otimes}$. Therefore, $(E[X])^{\otimes} \subseteq (E^{\otimes_0}[X])^{\otimes} \subseteq ((E[X])^{\otimes})^{\otimes} = (E[X])^{\otimes}$.

(3) Let $z \in E^{(\otimes_f)_0}$. Then there exists $F \in f(D[X])$ such that $F \subseteq E[X]$ and $z \in F^{\otimes} \cap K$. Let $I := c_p(F)$. Clearly, $I \in f(D)$, $I \subseteq E$, and $F \subseteq I[X] \subseteq E[X]$. Therefore, $z \in F^{\otimes} \cap K \subseteq (I[X])^{\otimes} \cap K = I^{\otimes_0}$, and so $z \in E^{(\otimes_0)_f}$. Conversely, if $z \in E^{(\otimes_0)_f}$, then $z \in I^{\otimes_0} = (I[X])^{\otimes} \cap K$ for some $I \in f(D)$, $I \subseteq E$. This implies that $z \in E^{(\otimes_f)_0}$.

(4) The “finite type part” is a particular case of (3). The “stable part” is a straightforward consequence of the definitions.

(5) is straightforward.

(6) Clearly, $(\tilde{\otimes})_0 \leq \tilde{\otimes}_0$, since $(\tilde{\otimes})_0 \leq \otimes_0$ by (5) and $(\tilde{\otimes})_0$ is a stable operation of finite type by statement (4). Let $a \in E^{\otimes_0}$ with $E \in \overline{\mathbf{F}}(D)$. Then, there exists a nonzero finitely generated ideal J of D such that $J^{\otimes_0} = D^{\otimes_0}$ and $aJ \subseteq E$. On the other hand, $(J[X])^{\otimes} = (J^{\otimes_0}[X])^{\otimes} = (D^{\otimes_0}[X])^{\otimes} = (D[X])^{\otimes}$ by (2). Since $J[X]$ is a nonzero finitely generated ideal of $D[X]$, $(J[X])^{\otimes} = (D[X])^{\otimes}$, and $aJ[X] \subseteq E[X]$, then $a \in (E[X])^{\otimes}$, and so $a \in (E[X])^{\otimes} \cap K = E^{\otimes_0}$.

(7) The statement for the d -operations is trivial. For the w -, t -, and v -operation, it is an easy consequence of the following equalities [19, Proposition 4.3] for all fractional ideals $E \in \mathbf{F}(D)$ (see also [32]).

$$(E[X])^{v_{D[X]}} = E^{v_D}[X], \quad (E[X])^{t_{D[X]}} = E^{t_D}[X], \quad (E[X])^{w_{D[X]}} = E^{w_D}[X].$$

(Note that if $E \in \overline{\mathbf{F}}(D) \setminus \mathbf{F}(D)$, then $E[X] \in \overline{\mathbf{F}}(D[X]) \setminus \mathbf{F}(D[X])$, and so $E^{v_D}[X] = K[X]$ and $(E[X])^{v_{D[X]}} = K(X)$; however, $(E[X])^{v_{D[X]}} \cap K = K = E^{v_D}$.)

We want to prove next that

$$(E[X])^{b_{D[X]}} = E^{b_D}[X], \quad \text{for all } E \in \overline{\mathbf{F}}(D).$$

We use the fact that $E^{b_D} = \bigcup \{(EI : I) \mid I \in \mathbf{f}(D)\}$ (respectively, $(E[X])^{b_{D[X]}} = \bigcup \{(E[X]F : F) \mid F \in \mathbf{f}(D[X])\}$) (see [36, p. 349] and [14, Section 19.3]). Let $z \in E^{b_D}[X] (\subseteq K[X])$. Then $z \in (EI : I)[X] = (E[X]I[X] : I[X])$ for some $I \in \mathbf{f}(D)$, and so, in particular, $z \in \bigcup \{(E[X]F : F) \mid F \in \mathbf{f}(D[X])\}$. Conversely, let $z \in \bigcup \{(E[X]F : F) \mid F \in \mathbf{f}(D[X])\} (\subseteq K[X])$. Then $zF \subseteq E[X]F$, and so (by [12, Theorem 28.1]) there exists a positive integer m such that $c_D(z)(c_D(z)^m c_D(F)) = c_D(z)^{m+1} c_D(F) = c_D(z)^m c_D(zF) \subseteq c_D(z)^m c_D(E[X]F) \subseteq c_D(z)^m c_D(E[X]) c_D(F) = E(c_D(z)^m c_D(F))$, where $c_D(z)^m c_D(F) \in \mathbf{f}(D)$. Therefore, $c_D(z) \subseteq E^{b_D}$, and thus $z \in c_D(z)[X] \subseteq E^{b_D}[X]$. \square

Remark 6. Note that the equality $(E[X])^{b_{D[X]}} = E^{b_D}[X]$, for all $E \in \overline{\mathbf{F}}(D)$ proved in (7) of Lemma 5, is equivalent to each of the following equalities:

$$\begin{aligned} E[X] \text{Kr}(D[X], b_{D[X]}) \cap K(X) &= (E \text{Kr}(D, b_D) \cap K)[X] \\ &= E \text{Kr}(D, b_D) \cap K[X], \\ &= \bigcap \{E[X]W \mid W \text{ valuation overring of } D[X]\} \\ &= \bigcap \{EV \mid V \text{ valuation overring of } D\}[X]. \end{aligned}$$

Remark 7. Given an arbitrary multiplicative subset \mathcal{S} of $D[X]$, Chang and Fontana in [3] investigated the map $E \mapsto ED[X]_{\mathcal{S}} \cap K$, defined for all $E \in \overline{\mathbf{F}}(D)$, showing that it gives rise to a semistar operation \star on D , having the properties that $D^{\star} = R := D[X]_{\mathcal{S}} \cap K$, and that R is t -linked to (D, \star) (i.e., for each nonzero finitely generated ideal I of D , $I^{\star} = D^{\star}$ implies $(IR)^{t_R} = R$ [5, Section 3]; or, equivalently, $R = R^{\tilde{\star}}$ [4, Lemma 2.9]). One of the main results of the article by Chang and

Fontana [3, Theorem 2.1] is recalled below, since it is strictly linked to the theme of the present work.

Note that, to a multiplicative subset \mathcal{S} of $D[X]$, we can associate the semistar operation $\otimes_{\mathcal{S}}$ on $D[X]$ defined by $A^{\otimes_{\mathcal{S}}} := A_{\mathcal{S}} = \bigcup \{(A : J) \mid J \text{ ideal of } D[X], J \cap \mathcal{S} \neq \emptyset\} = AD[X]_{\mathcal{S}}$, for all $A \in \overline{F}(D[X])$ [6, Proposition 2.10]. Therefore, by Lemma 5, we obtain immediately that the map $E \mapsto ED[X]_{\mathcal{S}} \cap K =: E^{\odot_{\mathcal{S}}}$, defined for all $E \in \overline{F}(D)$, gives rise to a semistar operation $\odot_{\mathcal{S}}$ on D coinciding with $(\otimes_{\mathcal{S}})_0$. Clearly, if $\overline{\mathcal{S}} := D[X] \setminus \bigcup \{Q \mid Q \in \text{Spec}(D[X]) \text{ and } Q \cap \mathcal{S} = \emptyset\}$ is the saturation of the multiplicative set \mathcal{S} , then $\otimes_{\mathcal{S}} = \otimes_{\overline{\mathcal{S}}}$ and so, in particular, $\odot_{\mathcal{S}} = \odot_{\overline{\mathcal{S}}}$.

In order to deepen our knowledge of the semistar operation $\odot_{\mathcal{S}}$, we need a definition of a stronger version of saturation. Set

$$\mathcal{S}^{\#} := D[X] \setminus \bigcup \{P[X] \mid P \in \text{Spec}(D) \text{ and } P[X] \cap \mathcal{S} = \emptyset\}.$$

It is clear that $\mathcal{S}^{\#}$ is a saturated multiplicative set of $D[X]$ and that $\mathcal{S}^{\#}$ contains the saturation of \mathcal{S} , i.e., $\mathcal{S}^{\#} \supseteq \overline{\mathcal{S}} \supseteq \mathcal{S}$. We call $\mathcal{S}^{\#}$ the *extended saturation* of \mathcal{S} in $D[X]$ and a multiplicative set \mathcal{S} of $D[X]$ is called *extended saturated* if $\mathcal{S} = \mathcal{S}^{\#}$. Clearly, in general, $\otimes_{\mathcal{S}^{\#}} \geq \otimes_{\mathcal{S}} (= \otimes_{\overline{\mathcal{S}}})$. However, it can be shown that $(\otimes_{\mathcal{S}^{\#}})_0 = (\otimes_{\mathcal{S}})_0$. For this, let

$$\Delta := \Delta(\mathcal{S}) := \{P \in \text{Spec}(D) \mid P[X] \cap \mathcal{S} = \emptyset\};$$

obviously, $\Delta(\mathcal{S}) = \Delta(\mathcal{S}^{\#})$. Let $\mathbf{V} := \mathbf{V}(\mathcal{S})$ be the set of the maximal elements of $\Delta(\mathcal{S})$. Then, by [3, Theorem 2.1], we have:

- (a) $\odot_{\mathcal{S}}$ is stable and of finite type, i.e., $\odot_{\mathcal{S}} = \widetilde{\odot_{\mathcal{S}}}$.
- (b) The extended saturation $\mathcal{S}^{\#}$ of \mathcal{S} coincides with $\mathcal{N}^{\odot_{\mathcal{S}}} := \{g \in D[X] \mid g \neq 0 \text{ and } c_D(g)^{\odot_{\mathcal{S}}} = D^{\odot_{\mathcal{S}}}\}$ and $\odot_{\mathcal{S}^{\#}} = \odot_{\mathcal{S}^{\#}}$.
- (c) If \mathcal{S} is extended saturated, then $\text{Na}(D, \odot_{\mathcal{S}}) = D[X]_{\mathcal{S}}$.
- (d) $\text{QMax}^{\odot_{\mathcal{S}}}(D) = \mathbf{V}(\mathcal{S})$. In particular, $\odot_{\mathcal{S}}$ coincides with the spectral semistar operation associated to $\mathbf{V}(\mathcal{S})$, i.e.,

$$E^{\odot_{\mathcal{S}}} = \bigcap \{ED_P \mid P \in \mathbf{V}(\mathcal{S})\}, \quad \text{for all } E \in \overline{F}(D).$$

- (e) $\odot_{\mathcal{S}}$ is a (semi)star operation on D if and only if $\mathcal{S} \subseteq \mathcal{N}^{v_D} := \{g \in D[X] \mid g \neq 0 \text{ and } c_D(g)^{v_D} = D\}$ or, equivalently, if and only if $D = \bigcap \{D_P \mid P \in \mathbf{V}(\mathcal{S})\}$.
- (f) The map $\mathcal{S} \mapsto \odot_{\mathcal{S}}$ establishes a 1-1 correspondence between the extended saturated multiplicative subsets of $D[X]$ (respectively, extended saturated multiplicative subsets of $D[X]$ contained in \mathcal{N}^{v_D}) and the set of the stable semistar (respectively, (semi)star) operations of finite type on D .
- (g) Let \mathcal{S} be an extended saturated multiplicative set of $D[X]$. Then, $\text{Na}(D, v_D) = D[X]_{\mathcal{S}}$ if and only if $\mathcal{S} = \mathcal{N}^{v_D}$.
- (h) Let $R := D^{\odot_{\mathcal{S}}}$ and let $\iota : D \rightarrow R$ be the canonical embedding. The overring R is t -linked to $(D, \odot_{\mathcal{S}})$ and $\mathcal{S} \subseteq \mathcal{N}^{v_R} := \{g \in R[X] \mid g \neq 0 \text{ and } c_R(g)^{v_R} = R\}$ (i.e., $(\odot_{\mathcal{S}})_\iota$ is a (semi)star operation on R). Moreover $(\odot_{\mathcal{S}})_\iota = w_R$ if and only if the extended saturation $\mathcal{S}^{\#_R} := R[X] \setminus \bigcup \{Q[X] \mid Q \in \text{Spec}(R) \text{ and } Q[X] \cap \mathcal{S} = \emptyset\}$ of the multiplicative set \mathcal{S} in $R[X]$ coincides with \mathcal{N}^{v_R} .

From Remark 7(f), we deduce that each semistar operation \star on D which is stable and of finite type is equal to $(\otimes_{\mathcal{F}})_0 = \odot_{\mathcal{F}}$ for a unique extended saturated multiplicative set \mathcal{S} of $D[X]$.

Corollary 8. *Let \star be a finite type stable semistar operation on an integral domain D with field of quotients K , and let X be an indeterminate over K . Let $\mathcal{S}(\star) := \mathcal{N}^{\star} := \{g \in D[X] \mid c_D(g)^{\star} = D^{\star}\}$. Then:*

- (1) $\mathcal{S}(\star)$ is an extended saturated multiplicative set of $D[X]$ and, more precisely, $\mathcal{S}(\star) = D[X] \setminus \bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\}$ with $\mathcal{S}(\star)^{\sharp} = D[X] \setminus \bigcup \{P[X] \mid P \in \text{Spec}(D) \text{ and } P[X] \cap \mathcal{S}(\star) = \emptyset\} = \mathcal{S}(\star)$;
- (2) $\star = \odot_{\mathcal{S}(\star)} = (\otimes_{\mathcal{S}(\star)})_0$ and $\mathcal{S}(\star)$ is unique among the extended saturated multiplicative set \mathcal{S} of $D[X]$ for which $\odot_{\mathcal{S}} = \star$.

Proof. (1) Clearly, $\mathcal{S}(\star) = D[X] \setminus \bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\}$ since, for $0 \neq g \in D[X]$, $c_D(g)^{\star} = c_D(g)^{\star_f} = D^{\star}$ if and only if $c_D(g) \not\subseteq Q$ for all $Q \in \text{QMax}^{\star_f}(D)$ and, for each prime ideal Q of D , $c_D(g) \not\subseteq Q$ if and only if $g \notin Q[X]$. Moreover, clearly, $\bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\} \subseteq \bigcup \{P[X] \mid P \in \text{Spec}(D) \text{ and } P[X] \cap \mathcal{S}(\star) = \emptyset\}$. On the other hand, if $P \in \text{Spec}(D)$ and $P[X] \cap \mathcal{S}(\star) = \emptyset$, this means that $P[X] \subseteq \bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\}$ and so $\bigcup \{P[X] \mid P \in \text{Spec}(D) \text{ and } P[X] \cap \mathcal{S}(\star) = \emptyset\} \subseteq \bigcup \{Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\}$.

(2) Note that, by assumption, $\star = \tilde{\star}$ and so, for each $E \in \overline{F}(D)$, $E^{\star} = E\text{Na}(D, \star) \cap K = ED[X]_{\mathcal{N}^{\star}} \cap K = ED[X]_{\mathcal{S}(\star)} \cap K$ [10, Proposition 3.4]. Therefore, if $\otimes_{\mathcal{S}(\star)}$ is the finite type stable semistar operation on $D[X]$ defined by the flat overring $D[X]_{\mathcal{S}(\star)}$, i.e., $A^{\otimes_{\mathcal{S}(\star)}} := A_{\mathcal{S}(\star)} = AD[X]_{\mathcal{S}(\star)}$ for all $A \in \overline{F}(D[X])$, then $\star = (\otimes_{\mathcal{S}(\star)})_0 = \odot_{\mathcal{S}(\star)}$. The uniqueness follows from Remark 7(f). \square

Note that from $E^{\otimes_0} = (E[X])^{\otimes} \cap K$, by tensoring with the D -algebra $D[X]$, we have $E^{\otimes_0}[X] = (E[X])^{\otimes} \cap K[X]$, for all $E \in \overline{F}(D)$. Moreover, it may happen that $E^{\otimes_0}[X] \subsetneq (E[X])^{\otimes}$ for some $E \in \overline{F}(D)$. For instance, this happens if $E^{\otimes_0} = K$ and if $K[X]$ is not a \otimes -overring of $D[X]$ (i.e., if $K[X] \subsetneq K[X]^{\otimes}$). An explicit example is given by $\otimes = v_{D[X]}$; in this case $K^{(v_{D[X]})_0}[X] = K[X] \subsetneq (K[X])^{v_{D[X]}} = K(X)$. Another example (even in case of finite type stable semistar operations) is given next by using Corollary 8.

Example 9. Let P be a given nonzero prime ideal of an integral domain D . Let $\Delta := \{P\}$, and set $\star := \star_{\Delta}$, i.e., \star is the finite type stable semistar operation defined by $E^{\star} := ED_P$, for all $E \in \overline{F}(D)$. Clearly, $\text{QMax}^{\star_{\Delta}}(D) = \{P\}$. Thus, $\mathcal{S}(\star) := D[X] \setminus P[X]$ and $\star = \tilde{\star} = \odot_{\mathcal{S}(\star)} = (\otimes_{\mathcal{S}(\star)})_0$, by Corollary 8 (2). (Note that $A^{\otimes_{\mathcal{S}(\star)}} = AD[X]_{P[X]} = AD_P(X)$ for each $A \in \overline{F}(D[X])$.) On the other hand, for each $E \in \overline{F}(D)$, $E^{\star}[X] = ED_P[X] \subsetneq ED[X]_{P[X]} = E[X]D_P(X) = (E[X])^{\otimes_{\mathcal{S}(\star)}}$ (even if $E^{\star}[X] = ED_P[X] = (ED_P(X) \cap K)[X] = E[X]D_P(X) \cap K[X] = (E[X])^{\otimes_{\mathcal{S}(\star)}} \cap K[X]$).

In order to better investigate this situation, we introduce the following definitions. A semistar operation \otimes on the polynomial domain $D[X]$ is called an *extension* (respectively, a *strict extension*) of a semistar operation \star defined on D if $E^{\star} = (E[X])^{\otimes} \cap K$ (respectively, $E^{\star}[X] = (E[X])^{\otimes}$) for all $E \in \overline{F}(D)$. Clearly, a strict

extension is an extension. By Lemma 5, a semistar operation \otimes on $D[X]$ is an extension of $\star := \otimes_0$.

Given two semistar operations \otimes' and \otimes'' on the polynomial domain $D[X]$, we say that they are *equivalent over D* , for short $\otimes' \sim \otimes''$, (respectively, *strictly equivalent over D* , for short $\otimes' \approx \otimes''$) if $(E[X])^{\otimes'} \cap K = (E[X])^{\otimes''} \cap K$ (respectively, $(E[X])^{\otimes'} = (E[X])^{\otimes''}$) for each $E \in \overline{F}(D)$.

Clearly, two extensions (respectively, strict extensions) \otimes' and \otimes'' on $D[X]$ of the same semistar operation defined on D are equivalent (respectively, strictly equivalent). In particular, we have

$$\otimes' \approx \otimes'' \Rightarrow \otimes' \sim \otimes'' \Leftrightarrow \otimes'_0 = \otimes''_0.$$

We will see that the converse of the first implication above does not hold in general. In order to construct some counterexamples, we need a deeper study of the problem of “raising” semistar operations from D to $D[X]$; i.e., given a semistar operation \star on D , finding all the semistar operations \otimes on $D[X]$ such that $\star = \otimes_0$.

Recall that, given a family of semistar operations $\{\star_\lambda \mid \lambda \in \Lambda\}$ on an integral domain D , the semistar operation $\wedge \star_\lambda$ on D is defined for all $E \in \overline{F}(D)$ by setting

$$E^{\wedge \star_\lambda} := \bigcap \{E^{\star_\lambda} \mid \lambda \in \Lambda\}.$$

The following statement is a straightforward consequence of the definitions.

Proposition 10.

- (1) Let \star be a semistar operation on an integral domain D . Given a family of semistar operations $\{\otimes_\lambda \mid \lambda \in \Lambda\}$ on $D[X]$ that are extensions (respectively, strict extensions) of \star , then $\wedge \otimes_\lambda$ is also an extension (respectively, a strict extension) of \star .
- (2) Given a family of semistar operations $\{\otimes_\lambda \mid \lambda \in \Lambda\}$ of $D[X]$, suppose that $\otimes_{\lambda'} \sim \otimes_{\lambda''}$ (respectively, $\otimes_{\lambda'} \approx \otimes_{\lambda''}$) for all $\lambda', \lambda'' \in \Lambda$, then $\wedge \otimes_\lambda$ is equivalent (respectively, strictly equivalent) to $\otimes_{\bar{\lambda}}$ for each $\bar{\lambda} \in \Lambda$.

From the previous proposition, we deduce that, if a semistar operation on D admits an extension (respectively, a strict extension) to $D[X]$, then it admits a unique minimal extension (respectively, a unique minimal strict extension).

At this point, it is natural to ask the following questions:

(Q1) Given a semistar operation \star defined on D , is it possible to find “in a canonical way” an extension (respectively, a strict extension) of \star on $D[X]$?

(Q2) Given an extension \otimes on $D[X]$ of a semistar operation \star defined on D . Is it possible to define a strict extension \otimes' on $D[X]$ of \star (and thus $\otimes' \sim \otimes$)? (In the statement of the previous question, we do not require that $\otimes' \approx \otimes$, since this condition would imply that the extension \otimes on $D[X]$ was already a strict extension of \star .)

In the remaining part of this paper, we start the investigation of questions (Q1) and (Q2), by considering semistar operations on D defined by families of overrings.

In this particular, but rather important setting, we will provide positive answers to both questions.

Let $\mathcal{T} := \{T_\lambda \mid \lambda \in \Lambda\}$ be a nonempty set of overrings of D , and let $E^{\wedge_{\mathcal{T}}} := \bigcap_{\lambda} ET_\lambda$ for each $E \in \overline{F}(D)$. Then $\wedge_{\mathcal{T}}$ is a semistar operation on D , and $\wedge_{\mathcal{T}}$ is (semi)star if and only if $D = \bigcap_{\lambda} T_\lambda$. It is easy to see that, for each $E \in \overline{F}(D)$ and for each $\lambda \in \Lambda$,

$$E^{\wedge_{\mathcal{T}}} T_\lambda = ET_\lambda,$$

(see [1, Theorem 2] for further details in the star operation case). If $\mathcal{T} = \{K\}$ (respectively, $\{D\}$) then obviously $\wedge_{\{K\}}$ (respectively, $\wedge_{\{D\}}$) is the trivial semistar operation $e_D (= \star_{\{K\}})$ (respectively, the identity (semi)star operation $d_D (= \star_{\{D\}})$). In case $\mathcal{T} = \emptyset$, we also set $\wedge_{\emptyset} := e_D$.

Note that, for each T_λ , $ET_\lambda = \bigcap \{(ET_\lambda)_M \mid M \in \text{Max}(T_\lambda)\} = \bigcap \{E(T_\lambda)_M \mid M \in \text{Max}(T_\lambda)\}$; hence $E^{\wedge_{\mathcal{T}}} = \bigcap_{\lambda} (\bigcap \{E(T_\lambda)_M \mid M \in \text{Max}(T_\lambda)\})$. If \mathcal{T} is nonempty, replacing the family $\mathcal{T} = \{T_\lambda \mid \lambda \in \Lambda\}$ with the family $\{(T_\lambda)_M \mid \lambda \in \Lambda, M \in \text{Max}(D_\lambda)\}$, without loss of generality, whenever convenient for the context, we can assume that each T_λ in the family of overrings \mathcal{T} is a quasi-local domain.

If \mathcal{T}' and \mathcal{T}'' are two families of overrings of D , then clearly

$$\wedge_{\mathcal{T}'} \wedge_{\mathcal{T}''} = \wedge_{\mathcal{T}' \cup \mathcal{T}''}.$$

Let $\mathcal{T} = \{T_\lambda \mid \lambda \in \Lambda\}$ be a family of overrings of an integral domain D with quotient field K . Let X be an indeterminate over K , and denote by $T_\lambda(X)$ the Nagata ring of T_λ . For each $A \in \overline{F}(D[X])$, we set

$$\begin{aligned} A^{(\wedge_{\mathcal{T}})} &:= \bigcap_{\lambda} AT_\lambda(X), \\ A^{\langle \wedge_{\mathcal{T}} \rangle} &:= A^{(\wedge_{\mathcal{T}})} \cap AK[X], \\ A^{[\wedge_{\mathcal{T}}]} &:= \bigcap_{\lambda} AT_\lambda[X]. \end{aligned}$$

Clearly, $A^{[\wedge_{\mathcal{T}}]} \subseteq A^{\langle \wedge_{\mathcal{T}} \rangle} \subseteq A^{(\wedge_{\mathcal{T}})}$ for all $A \in \overline{F}(D[X])$, and hence $[\wedge_{\mathcal{T}}] \leq \langle \wedge_{\mathcal{T}} \rangle \leq (\wedge_{\mathcal{T}})$. Moreover, if \mathcal{T} is nonempty, $(D[X])^{\langle \wedge_{\mathcal{T}} \rangle} \subseteq K[X]$, but $1/(1+X) \in \bigcap_{\lambda} T_\lambda(X) = (D[X])^{(\wedge_{\mathcal{T}})}$. Hence, $(D[X])^{\langle \wedge_{\mathcal{T}} \rangle} \subsetneq (D[X])^{(\wedge_{\mathcal{T}})}$, and so $\langle \wedge_{\mathcal{T}} \rangle \lneq (\wedge_{\mathcal{T}})$.

Proposition 11. *Let $\mathcal{T} = \{T_\lambda \mid \lambda \in \Lambda\}$ be a family of overrings of an integral domain D with quotient field K . Set $(\mathcal{T}) := \{T_\lambda(X) \mid \lambda \in \Lambda\}$, $\langle \mathcal{T} \rangle := \{T_\lambda(X) \mid \lambda \in \Lambda\} \cup \{K[X]\}$, and $[\mathcal{T}] := \{T_\lambda[X] \mid \lambda \in \Lambda\}$. Then:*

(1) $(\wedge_{\mathcal{T}})$, $\langle \wedge_{\mathcal{T}} \rangle$, and $[\wedge_{\mathcal{T}}]$ are semistar operations on $D[X]$ and, more precisely,

$$(\wedge_{\mathcal{T}}) = \wedge_{(\mathcal{T})}, \quad \langle \wedge_{\mathcal{T}} \rangle = \wedge_{\langle \mathcal{T} \rangle}, \quad \text{and} \quad [\wedge_{\mathcal{T}}] = \wedge_{[\mathcal{T}]}.$$

Moreover, if we consider the set $\mathcal{U} := \{K[X]\}$ consisting of the unique overring $K[X]$ of $D[X]$, then

$$\langle \wedge_{\mathcal{T}} \rangle = (\wedge_{\mathcal{T}}) \wedge \wedge_{\mathcal{U}} (= \wedge_{(\mathcal{T})} \wedge \wedge_{\mathcal{U}}).$$

(2) The following are equivalent:

- (i) $\wedge_{\mathcal{T}}$ is a (semi)star operation on D ;
- (ii) $\langle \wedge_{\mathcal{T}} \rangle$ is a (semi)star operation on $D[X]$;
- (iii) $[\wedge_{\mathcal{T}}]$ is a (semi)star operation on $D[X]$.

(3) If \mathcal{T} is not empty and $\mathcal{T} \neq \{K\}$, then $[\wedge_{\mathcal{T}}] \leq \langle \wedge_{\mathcal{T}} \rangle$.

(4) For each $E \in \overline{F}(D)$,

$$(E[X])^{[\wedge_{\mathcal{T}}]} = E^{\wedge_{\mathcal{T}}}[X] = (E[X])^{\langle \wedge_{\mathcal{T}} \rangle} \quad \text{and} \quad (E[X])^{\langle \wedge_{\mathcal{T}} \rangle} = E^{\wedge_{\mathcal{T}}}(X).$$

(5) For each $E \in \overline{F}(D)$,

$$E^{\wedge_{\mathcal{T}}} = (E[X])^{[\wedge_{\mathcal{T}}]} \cap K = (E[X])^{\langle \wedge_{\mathcal{T}} \rangle} \cap K = (E[X])^{(\wedge_{\mathcal{T}})} \cap K.$$

(6) If \mathcal{T} is a finite family of overrings of D , then $\wedge_{\mathcal{T}}$ is a semistar operation of finite type on D .

(7) If each overring $T \in \mathcal{T}$ is a flat overring of D , then $\wedge_{\mathcal{T}}$ is a stable semistar operation on D .

Proof. (1) is obvious and (2) is straightforward, since it is easy to see that $\bigcap \{T_{\lambda}[X] \mid \lambda \in \Lambda\} = (\bigcap \{T_{\lambda} \mid \lambda \in \Lambda\})[X]$ and $\bigcap \{T_{\lambda}(X) \mid \lambda \in \Lambda\} = (\bigcap \{T_{\lambda} \mid \lambda \in \Lambda\})(X)$.

(3) Without loss of generality, we can assume that T_{λ} is local with nonzero maximal ideal M_{λ} and we can take the (maximal) ideal $(M_{\lambda}, 1 + X)$ of $T_{\lambda}[X]$. Set $Q := (M_{\lambda}, 1 + X) \cap D[X]$. Then, $Q^{[\wedge_{\mathcal{T}}]} \subsetneq Q^{\langle \wedge_{\mathcal{T}} \rangle}$ (for $Q^{[\wedge_{\mathcal{T}}]} \cap D[X] \subseteq (M_{\lambda}, 1 + X) \cap D[X] = Q \subsetneq D[X]$ and, on the other hand, since $M_{\lambda} \neq (0)$ and $Q^{\langle \wedge_{\mathcal{T}} \rangle} = D[X]^{\langle \wedge_{\mathcal{T}} \rangle}$, then $Q^{\langle \wedge_{\mathcal{T}} \rangle} \cap D[X] = Q^{\langle \wedge_{\mathcal{T}} \rangle} \cap QK[X] \cap D[X] = D[X]^{\langle \wedge_{\mathcal{T}} \rangle} \cap D[X] = D[X]$).

(4) is straightforward, and (5) is a trivial consequence of (4).

(6) Clearly, for each $T \in \mathcal{T}$, the operation $\wedge_{\{T\}}$ (defined by $A^{\wedge_{\{T\}}} := AT$, for all $A \in \overline{F}(D)$, i.e., $\wedge_{\{T\}} = \star_{\{T\}}$) is a semistar operation of finite type on D . Let $\mathcal{T} := \{T_1, T_2, \dots, T_n\}$, and let $z \in A^{\wedge_{\mathcal{T}}}$. Then, $z \in F_i T_i$, for some $F_i \subseteq A$, $F_i \in \mathbf{f}(D)$ and for each $1 \leq i \leq n$. Set $F := F_1 + F_2 + \dots + F_n$; clearly, $F \in \mathbf{f}(D)$; $F \subseteq A$, and $z \in F T_i$ for all i , thus $z \in F^{\wedge_{\mathcal{T}}}$.

(7) Let $A, B \in \overline{F}(D)$. Since T_{λ} is flat, $(A \cap B)T_{\lambda} = AT_{\lambda} \cap BT_{\lambda}$ [2, Chapter 1, §2, N. 6]. The conclusion is now straightforward. \square

From Proposition 11, we deduce immediately the following corollary.

Corollary 12. *With the notation of Proposition 11, assume that \mathcal{T} is nonempty and $\mathcal{T} \neq \{K\}$. Then we have*

- (1) $[\wedge_{\mathcal{T}}]$, $\langle \wedge_{\mathcal{T}} \rangle$, and $(\wedge_{\mathcal{T}})$ (respectively, $[\wedge_{\mathcal{T}}]$ and $\langle \wedge_{\mathcal{T}} \rangle$) are distinct extensions (respectively, distinct strict extensions) of $\wedge_{\mathcal{T}}$;
- (2) $[\wedge_{\mathcal{T}}] \sim \langle \wedge_{\mathcal{T}} \rangle \sim (\wedge_{\mathcal{T}})$ and, moreover, $[\wedge_{\mathcal{T}}] \approx \langle \wedge_{\mathcal{T}} \rangle$, but neither $[\wedge_{\mathcal{T}}]$ nor $\langle \wedge_{\mathcal{T}} \rangle$ are strictly equivalent to $(\wedge_{\mathcal{T}})$.

Example 13. Let $\mathcal{W} := \{W_\lambda \mid \lambda \in \Lambda\}$ be a family of valuation overrings of D , and let $\wedge_{\mathcal{W}}$ be the ab semistar operation on D defined by the family of valuation overrings \mathcal{W} of D (i.e., $E^{\wedge_{\mathcal{W}}} := \bigcap \{EW \mid W \in \mathcal{W}\}$ for all $E \in \overline{F}(D)$) [9, Proposition 3.7(1)]. In this case,

- (a) $(\wedge_{\mathcal{W}})$ is an ab semistar operation on $D[X]$ defined by the family of valuation overrings $\mathcal{W}(X) := \{W_\lambda(X) \mid \lambda \in \Lambda\}$ of $D[X]$ and (by [9, Corollary 3.8]) for each $A \in \overline{F}(D[X])$,

$$A^{(\wedge_{\mathcal{W}})} = \bigcap_{\lambda} AW_\lambda(X);$$

- (b) For each $E \in \overline{F}(D)$,

$$\begin{aligned} E^{(\wedge_{\mathcal{W}})_0} &= (E[X])^{(\wedge_{\mathcal{W}})} \cap K = \left(\bigcap_{\lambda} EW_\lambda(X) \right) \cap K \\ &= \bigcap_{\lambda} (EW_\lambda(X) \cap K) = \bigcap_{\lambda} EW_\lambda \\ &= E^{\wedge_{\mathcal{W}}}; \end{aligned}$$

- (c) For each $F \in f(D)$,

$$F^{(\wedge_{\mathcal{W}})_0} = F^{\wedge_{\mathcal{W}}} = F\text{Kr}(D, \wedge_{\mathcal{W}}) \cap K.$$

In particular, by (b), Lemma 5(2) and [11, Proposition 9]

$$(\wedge_{\mathcal{W}})_{0,f} = (\wedge_{\mathcal{W}})_{f,0} = (\wedge_{\mathcal{W}})_{a,0} = \wedge_{\mathcal{W},a} = \wedge_{\mathcal{W},f}$$

(where $(\wedge_{\mathcal{W}})_{0,f}$ (respectively, $(\wedge_{\mathcal{W}})_{f,0}$; $(\wedge_{\mathcal{W}})_{a,0}$; $\wedge_{\mathcal{W},a}$; $\wedge_{\mathcal{W},f}$) denotes the semistar operation of finite type on D associated to $(\wedge_{\mathcal{W}})_0$ (respectively, the semistar operation on D canonically induced by $(\wedge_{\mathcal{W}})_f$; the semistar operation on D canonically induced by $(\wedge_{\mathcal{W}})_a$; the ab semistar operation on D associated to $\wedge_{\mathcal{W}}$; the semistar operation of finite type on D associated to $\wedge_{\mathcal{W}}$).

Example 14.

- (1) The identity (semi)star operation d_D on an integral domain D , is defined by the family of a single overring $\mathcal{D} := \{D\}$ of D , i.e., $d_D = \wedge_{\mathcal{D}}$. Set

$$[d_D] := [\wedge_{\mathcal{D}}], \quad \langle d_D \rangle := \langle \wedge_{\mathcal{D}} \rangle, \quad (d_D) := (\wedge_{\mathcal{D}}).$$

Clearly, if D is not a field, $d_{D[X]} = [d_D] \leq \langle d_D \rangle \leq (d_D)$ and $\langle d_D \rangle$ (and $[d_D]$) is a (semi)star operation on $D[X]$, but in general (d_D) is not a (semi)star operation on $D[X]$. Moreover, $\langle d_D \rangle$, (d_D) (and $[d_D]$) are stable semistar operations of finite type, since $\langle d_D \rangle$ is defined by the two flat overrings $D(X)$ and $K[X]$ of $D[X]$ and (d_D) is defined by a unique flat overring $D(X)$ of $D[X]$ (Proposition 11(6) and (7)).

- (2) As observed above, the trivial semistar operation e_D on an integral domain D , with quotient field K , is defined by the family of a single overring $\mathcal{K} := \{K\}$ of D , i.e., $e_D = \wedge_{\mathcal{K}}$. Set

$$[e_D] := [\wedge_{\mathcal{K}}], \quad \langle e_D \rangle := \langle \wedge_{\mathcal{K}} \rangle, \quad (e_D) := (\wedge_{\mathcal{K}}).$$

Clearly, $[e_D] = \langle e_D \rangle \leq (e_D) = e_{D[X]}$, where $[e_D](= \langle e_D \rangle)$ is the stable semistar operation of finite type on $D[X]$ defined by the flat overring $K[X]$, i.e., $[e_D] = \langle e_D \rangle = \wedge_{\{K[X]\}} (= \star_{\{K[X]\}})$.

We study now the important case in which the family of valuation overrings \mathcal{W} of D coincides with the family of all valuation overrings of D .

Proposition 15. *Let \mathcal{V} be the family of all valuation overrings of an integral domain D with quotient field K . Note that $\wedge_{\mathcal{V}}$ coincides with b_D (the b -operation on D ; see Section 1). Set*

$$[b_D] := [\wedge_{\mathcal{V}}], \quad \langle b_D \rangle := \langle \wedge_{\mathcal{V}} \rangle, \quad (b_D) := (\wedge_{\mathcal{V}}).$$

- (1) $[b_D]$, $\langle b_D \rangle$, and (b_D) are semistar operations on $D[X]$ with $[b_D] \leq \langle b_D \rangle \leq (b_D)$. If D is integrally closed, then $[b_D]$ and $\langle b_D \rangle$ are (semi)star operations on $D[X]$. In general, (b_D) is not a (semi)star operation on $D[X]$ even if D is integrally closed (or, equivalently, even if b_D is a (semi)star operation on D).
- (2) If $D \neq K$, i.e., if D has at least one nontrivial valuation overring, then $[b_D]$, $\langle b_D \rangle$, and (b_D) (respectively, $[b_D]$ and $\langle b_D \rangle$) are distinct extensions (respectively, distinct strict extensions) of b_D .
- (3) $\langle b_D \rangle$ and (b_D) are **ab** semistar operations such that

$$[b_D] \leq [b_D]_a = b_{D[X]} \leq \langle b_D \rangle \leq (b_D)$$

and, in general, $[b_D]$ is not an **eab** semistar operation.

Proof. (1) and (2) follow from Proposition 11((1), (2), and (3)) and Corollary 12.

(3) Clearly, (b_D) is an **ab** semistar operation on $D[X]$, since it is defined by the family of valuation overrings $\mathcal{V}(X) := \{V(X) \mid V \in \mathcal{V}\}$.

Moreover, if $\mathcal{U} := \{K[X]\}$ and $\mathcal{U}' := \{K[X]_M \mid M \in \text{Max}(K[X])\}$, then clearly, $\wedge_{\mathcal{U}} = \wedge_{\mathcal{U}'}$. Therefore, $\langle b_D \rangle = \langle \wedge_{\mathcal{V}} \rangle = (\wedge_{\mathcal{V}}) \wedge \wedge_{\mathcal{U}} = \wedge_{\mathcal{V}(X)} \wedge \wedge_{\mathcal{U}'}$ and, hence $\langle b_D \rangle$ is also an **ab** semistar operation on $D[X]$.

Note that $b_{D[X]} \leq \langle b_D \rangle$ because $\mathcal{V}(X) \cup \mathcal{U}'$ is a subset of the family of all valuation overrings of $D[X]$.

Let W be a valuation overring of $D[X]$ with maximal ideal N . Two cases are possible. If $W \cap K = K$, then $K[X] \subseteq W$, and hence $W = K[X]_M$ for some maximal ideal M of $K[X]$. Next, if $W \cap K \subsetneq K$, then $V := W \cap K$ is a valuation overring of D with nonzero maximal ideal \mathfrak{n} , and so W is a valuation overring of $V[X]$. If $W \neq V(X) = V[X]_{\mathfrak{n}[X]}$ the maximal ideal N of W must contract on a maximal ideal of $V[X]$ upper to the maximal ideal \mathfrak{n} of V , i.e., $N \cap V[X] \supsetneq \mathfrak{n}[X]$. Therefore, we

have necessarily that $V[X] \subset W \subseteq V(X)$. From the previous observations, we easily deduce that $[b_D]_a = b_{D[X]}$.

We next construct an integral domain D such that $[b_D]$ is not an **eab** semistar operation. Let $D := \mathbb{R} + T\mathbb{C}[[T]]$, i.e., D is a pseudo-valuation domain with canonically associated valuation overring $V := \mathbb{C}[[T]]$ and quotient field $K := \mathbb{C}((T))$. Since $\mathbb{R} \subset \mathbb{C}$ is a finite field extension the valuation overrings of D are just V and K , thus it is straightforward to see that that $d_D \leq b_D = \wedge_{\{V\}}$ and

$$d_{D[X]} \leq [b_D] = \wedge_{\{V[X], K[X]\}} \leq b_{D[X]} \leq \langle b_D \rangle = \wedge_{\{V(X), K[X]\}} \leq (b_D) = \wedge_{\{V(X)\}}$$

(Proposition 11(3)). Moreover, $[b_D]$ is not an **eab** semistar operation on $D[X]$, because if $[b_D](= \wedge_{\{V[X], K[X]\}} = \star_{\{V[X]\}})$ was an **eab** semistar operation on $D[X]$, since it is of finite type, then $b_{D[X]} = (d_{D[X]})_a \leq [b_D] \leq b_{D[X]}$, i.e., $\star_{\{V[X]\}} = [b_D] = b_{D[X]}$, which is a contradiction since $V[X]$ is not a Prüfer domain. \square

In the next result, we provide another application of Proposition 11.

Proposition 16. *Let \star be a semistar operation of an integral domain D with quotient field K and let X be an indeterminate over K . Set $\mathcal{M} := \mathcal{M}(\star) := \{D_Q \mid Q \in Q\text{Max}^{\star'}(D)\}$. It is well known that, in this case, $\wedge_{\mathcal{M}}$ coincides with $\tilde{\star}$, the stable semistar operation of finite type associated to \star . Set*

$$(\tilde{\star}) := (\wedge_{\mathcal{M}}), \quad \langle \tilde{\star} \rangle := (\wedge_{\mathcal{M}}), \quad \text{and} \quad [\tilde{\star}] := [\wedge_{\mathcal{M}}].$$

(1) *For each $A \in \overline{F}(D[X])$,*

$$A^{[\tilde{\star}]} = \bigcap \{AD_Q[X] \mid Q \in Q\text{Max}^{\star'}(D)\},$$

$$A^{(\tilde{\star})} = ANa(D, \star) \cap AK[X], \quad \text{and} \quad A^{(\tilde{\star})} = ANa(D, \star).$$

(2) *$[\tilde{\star}]$, $\langle \tilde{\star} \rangle$, and $(\tilde{\star})$ are stable semistar operations of $D[X]$; moreover, $\langle \tilde{\star} \rangle$ and $(\tilde{\star})$ are also of finite type. Therefore,*

$$\langle \tilde{\star} \rangle = \langle \tilde{\star} \rangle_f = \widetilde{\langle \tilde{\star} \rangle}, \quad (\tilde{\star}) = (\tilde{\star})_f = \widetilde{(\tilde{\star})}.$$

(3) *For each $E \in \overline{F}(D)$,*

$$(ED[X])^{[\tilde{\star}]} = E^{\star}[X] = (ED[X])^{(\tilde{\star})} \quad \text{and} \quad (ED[X])^{(\tilde{\star})} = ENa(D, \star);$$

and so

$$E^{\star} = (ED[X])^{[\tilde{\star}]} \cap K = (ED[X])^{(\tilde{\star})} \cap K = (ED[X])^{(\tilde{\star})} \cap K.$$

Proof. (1) and (2). Note that the semistar operation $[\tilde{\star}]$ (respectively, $\langle \tilde{\star} \rangle$; $(\tilde{\star})$) on $D[X]$ is defined by the family of flat overrings $\{D_Q[X] \mid Q \in Q\text{Max}^{\star'}(D)\}$ (respectively, $\{D_Q(X) \mid Q \in Q\text{Max}^{\star'}(D)\} \cup \{K[X]\}$; $\{D_Q(X) \mid Q \in Q\text{Max}^{\star'}(D)\}$) of $D[X]$ and so is a stable semistar operation of $D[X]$. The first equality in (1) is a transcription of the definition. The last two equalities in (1) are consequences of

the fact that $\bigcap \{AD_Q(X) \mid Q \in \text{QMax}^{*f}(D)\} = \text{ANa}(D, \star)$ (Proposition 2(2) or [10, Proposition 3.1(3)]). Finally, it is easy to see that $(\tilde{\star})$ (respectively, $\langle \tilde{\star} \rangle$) is a (stable) semistar operation of finite type, since it is defined by a unique flat overring of $D[X]$, i.e., $\text{Na}(D, \star)$ (respectively, by two flat overrings of $D[X]$, i.e., $\text{Na}(D, \star)$ and $K[X]$) Proposition 11 (6) and (7).

(3) is an application of Proposition 11 (4) and (5). \square

It is natural to ask if, eventually, $[\tilde{\star}] \neq [\tilde{\star}]_f$. The answer to this question is negative (i.e., $[\tilde{\star}]$ is also a semistar operation of finite type (and stable)). In order to show this fact, we deepen the study of the semistar operation $[\tilde{\star}]$ defined on $D[X]$.

Proposition 17. *Let D , X , \star and $[\tilde{\star}]$ be as in Proposition 16. Then, for all $A \in \overline{F}(D[X])$,*

$$\begin{aligned} A^{[\tilde{\star}]} &= \bigcup \{(A : F) \mid F \in \mathbf{f}(D) \text{ and } F^{\tilde{\star}} = D^{\tilde{\star}}\} \\ &= \bigcup \{(A : H) \mid H \in \mathbf{f}(D), H \subseteq D, \text{ and } H^{\tilde{\star}} = D^{\tilde{\star}}\}. \end{aligned}$$

In particular, $[\tilde{\star}]$ is a (stable) semistar operation of finite type on $D[X]$ and so $[\tilde{\star}] = [\tilde{\star}]_f = \widetilde{[\tilde{\star}]}$.

Proof. Set $A_1 := \bigcup \{(A : F) \mid F \in \mathbf{f}(D) \text{ and } F^{\tilde{\star}} = D^{\tilde{\star}}\}$ and $A_2 := \bigcup \{(A : H) \mid H \in \mathbf{f}(D), H \subseteq D, \text{ and } H^{\tilde{\star}} = D^{\tilde{\star}}\}$. Clearly, $A_2 \subseteq A_1$.

We start by showing that $A_1 \subseteq A^{[\tilde{\star}]}$. Note that $F^{\tilde{\star}} = D^{\tilde{\star}}$ if and only if $FD_Q = D_Q$ for all $Q \in \text{QMax}^{*f}(D) = \text{QMax}^{\tilde{\star}}(D)$. If $z \in A_1$, then $zF \subseteq A$ for some $F \in \mathbf{f}(D)$ and $F^{\tilde{\star}} = D^{\tilde{\star}}$. Hence, $z \in zD_Q[X] = zFD_Q[X] \subseteq AD_Q[X]$ for all $Q \in \text{QMax}^{*f}(D)$ and so $z \in A^{[\tilde{\star}]}$.

Now, we show that $A^{[\tilde{\star}]} \subseteq A_2$. Let $z \in A^{[\tilde{\star}]} = \bigcap \{AD_Q[X] \mid Q \in \text{QMax}^{*f}(D)\}$ and set $I := \{d \in D \mid dz \in A\}$. It is easy to see that I is an ideal of D (depending on A and z) and, moreover, $I \not\subseteq Q$, i.e., $ID_Q = D_Q$ for all $Q \in \text{QMax}^{*f}(D)$. Since $I^{\tilde{\star}} = D^{\tilde{\star}}$, we can find $H \in \mathbf{f}(D)$ with $H \subseteq I$ and $H^{\tilde{\star}} = D^{\tilde{\star}}$. Therefore, $zH \subseteq zI \subseteq A$, and so $z \in A_2$.

The last statement follows easily from Proposition 16(2) and from the fact that, if $z \in A^{[\tilde{\star}]} = \bigcup \{(A : F) \mid F \in \mathbf{f}(D) \text{ and } F^{\tilde{\star}} = D^{\tilde{\star}}\}$, then $zF_0 =: G_0 \subseteq A$ for some $F_0 \in \mathbf{f}(D)$, with $F_0^{\tilde{\star}} = D^{\tilde{\star}}$, and so $G_0 \in \mathbf{f}(D)$, and $z \in (G_0 : F_0) \subseteq G_0^{[\tilde{\star}]} \subseteq A^{[\tilde{\star}]_f}$. \square

From Corollaries 8 and 12 and from Propositions 16 and 17, we easily obtain the following corollary.

Corollary 18. *Let \star , $\tilde{\star}$, D , K , X , $[\tilde{\star}]$, $\langle \tilde{\star} \rangle$, and $(\tilde{\star})$ be as in Proposition 16. Let $\mathcal{S}(\star)$ and $\otimes_{\mathcal{S}(\star)}$ be as in Corollary 8. Assume that $D_Q \subsetneq K$ for some $Q \in \text{QMax}^{*f}(D)$.*

- (1) $[\tilde{\star}]$, $\langle \tilde{\star} \rangle$, and $(\tilde{\star})$ (respectively, $[\tilde{\star}]$, and $\langle \tilde{\star} \rangle$) are distinct finite type stable semistar extensions to $D[X]$ (respectively, distinct finite type stable semistar strict extensions) of $\tilde{\star}$. Moreover, $(\tilde{\star})$ is the unique finite type stable semistar extension to $D[X]$ of $\tilde{\star}$ defined by an extended saturated multiplicative set of $D[X]$, i.e., $(\tilde{\star}) = \otimes_{\mathcal{S}(\star)}$, where $\mathcal{S}(\star) = \mathcal{N}^{\star} := \{g \in D[X] \mid c_D(g)^{\star} = D^{\star}\}$.
- (2) $[\tilde{\star}] \sim \langle \tilde{\star} \rangle \sim (\tilde{\star})$ and, moreover, $[\tilde{\star}] \approx \langle \tilde{\star} \rangle$, but neither $[\tilde{\star}]$ nor $\langle \tilde{\star} \rangle$ are strictly equivalent to $(\tilde{\star})$.

Remark 19. Picozza [30] has studied a different approach for the extension to the polynomial ring $D[X]$ of a semistar operation defined on an integral domain D .

First, he proves the following [30, Propositions 3.1 and 3.2].

(a) Let \mathcal{F} be a localizing system of ideals of D , and set:

$$\mathcal{F}[X] := \{J \text{ ideal of } D[X] \mid J \supseteq I[X] \text{ for some ideal } I \text{ of } \mathcal{F}\}.$$

(a.1) $\mathcal{F}[X]$ is a localizing system on $D[X]$,

(a.2) $\mathcal{F}[X] = \{J \text{ ideal of } D[X] \mid J \cap D \in \mathcal{F}\}$.

(a.3) If \mathcal{F} is a localizing system of finite type of D , then $\mathcal{F}[X]$ is a localizing system of finite type on $D[X]$.

Then, he uses some of the results by Fontana and Huckaba recalled in Proposition 1. More precisely, if \mathcal{F} is a localizing system on D , Picozza considers the semistar operation $\star_{\mathcal{F}[X]}$ on $D[X]$ canonically associated to the localizing system $\mathcal{F}[X]$ on $D[X]$ introduced in (a). In particular, if \mathcal{F} is the localizing system associated to a given semistar operation \star defined on D , i.e., $\mathcal{F} = \mathcal{F}^\star$, he considers the stable semistar operation on $D[X]$ associated to the localizing system $\mathcal{F}^\star[X]$. Set $\star[X] := \star_{\mathcal{F}[X]}$ (Picozza denotes by \star' this semistar operation on $D[X]$ [30, Theorem 3.3]).

We are now in a position to compare the semistar operations on the polynomial rings studied by Picozza and the semistar operation $[\tilde{\star}]$ introduced in Proposition 16. The following result improves [30, Proposition 3.4].

(b) Using the notation introduced above, then

$$\begin{aligned} [\tilde{\star}] &= \tilde{\star}[X] = \widetilde{\star[X]} \\ &= \star_f[X] = \star[X]_f. \end{aligned}$$

It is clear that $\tilde{\star}[X] = \star_f[X]$, since

$$\begin{aligned} \mathcal{F}^{\star_f} &= \{I \text{ ideal of } D \mid I^{\star_f} = D^{\star_f}\} \\ &= \{I \text{ ideal of } D \mid I \not\subseteq Q \text{ for all } Q \in \text{QMax}^{\star_f}(D)\} \\ &= \{I \text{ ideal of } D \mid I \not\subseteq Q \text{ for all } Q \in \text{QMax}^{\tilde{\star}}(D)\} \\ &= \{I \text{ ideal of } D \mid I^{\tilde{\star}} = D^{\tilde{\star}}\} \\ &= \mathcal{F}^{\tilde{\star}}. \end{aligned}$$

By Proposition 17, we know that $A^{[\tilde{\star}]} = \bigcup \{(A : H) \mid H \in f(D), H \subseteq D, \text{ and } H^{\tilde{\star}} = D^{\tilde{\star}}\}$ for all $A \in \overline{F}(D[X])$. On the other hand (by definition of a semistar operation associated to a localizing system Proposition 1(3) and by (a.2), we have $A^{\star[X]} = \bigcup \{(A : J) \mid J \text{ ideal of } D[X] \text{ such that } (J \cap D)^{\star} = D^{\star}\}$ for all $A \in \overline{F}(D[X])$. Therefore, if $z \in A^{[\tilde{\star}]}$, then $z \in (A : H) = (A : H[X])$ for some finitely generated ideal H of D such that $(H[X] \cap D)^{\tilde{\star}} = D^{\tilde{\star}}$, thus $z \in A^{\star[X]}$.

Conversely, let $z \in A^{\star[X]}$. Therefore, $z \in (A : J)$ for some ideal J of $D[X]$ such that $(J \cap D)^{\star} = D^{\star}$. In this situation, we can find a finitely generated ideal H in D such that $H \subseteq J \cap D$ and $H^{\star} = D^{\star}$. Since $H \subseteq J$, then $(A : J) \subseteq (A : H)$, thus $z \in (A : H)$ and so $z \in A^{[\tilde{\star}]}$.

From the previous results, we deduce:

- (c) $[d_D] = d_D[X] = d_{D[X]}$;
- (d) $[w_D] = w_D[X] = t_D[X] \leq w_{D[X]} \leq t_{D[X]}$.

The statement (c) is a straightforward consequence of (b), since $d_D = \widetilde{d}_D = d_{D,f}$ and the localizing system $\mathcal{F}^{d_D}[X] = \{D[X]\} = \mathcal{F}^{d_{D[X]}}$.

The equalities in (d) are obtained from (b) (and Proposition 16) by taking $\star = v_D$ (and so, $\tilde{\star} = w_D$ and $\star_f = t_D$). Moreover, it is always true that the w -operation is smaller than or equal to the t -operation. Finally, note that $w_D[X]$ (respectively, $w_{D[X]}$) is the stable (semi)star operation of finite type on $D[X]$ canonically associated to the localizing system $\mathcal{F}^{w_D}[X] = \{J \text{ ideal of } D[X] \mid J \supseteq I[X] \text{ with } I^{w_D} = D\}$ (respectively, $\mathcal{F}^{w_{D[X]}} = \{J \text{ ideal of } D[X] \mid J^{w_{D[X]}} = D[X]\}$) [6, Theorem 2.10(B)]. Since $I^{w_D} = D$ implies that $I^{w_D}[X] = D[X]$ and so also $I[X]^{w_{D[X]}} = D[X]$ (see the proof of Lemma 5(7)), then $\mathcal{F}^{w_D}[X] \subseteq \mathcal{F}^{w_{D[X]}}$. From this, we conclude that $w_D[X] \leq w_{D[X]}$.

Note that, in (d), it may happen that $[w_D] = w_D[X] \leq w_{D[X]}$ (e.g., by [30, Remark 2], let Q be a prime ideal of $D[X]$ not extended from D and such that $Q \cap D$ is a t_D -maximal (= w_D -maximal) ideal of D ; since $(Q \cap D)[X] \subsetneq Q$ and $(Q \cap D)[X]$ is a $t_{D[X]}$ -maximal (= $w_{D[X]}$ -maximal) ideal of $D[X]$ ([21, Proposition 1.1] and [8, Corollary 3.5(2)]), then Q is not a $w_{D[X]}$ -(maximal) ideal, but clearly Q is a $[w_D]$ -ideal).

Remark 20.

- (1) Using the techniques introduced in [3] and recalled in Remark 7, Sahandi [33, Theorem 2.1, Proposition 2.2 and Remark 2.3] has recently given another description of the stable semistar operation $[\tilde{\star}]$.

Let $D_1 := D[X]$, $K_1 := K(X)$, Y an indeterminate over K_1 , and consider the following subset of $\text{Spec}(D_1)$:

$$\Delta_1 := \Delta_{1,\star} := \{Q_1 \in \text{Spec}(D_1) \mid \text{either } Q_1 \cap D = (0) \text{ or } (Q_1 \cap D)^{\star_f} \subsetneq D^{\star_f}\}.$$

Set

$$\mathcal{S}_1^\# := \mathcal{S}_{1,\star}^\# := \mathcal{S}^\#(\Delta_{1,\star}) := D_1[Y] \setminus \left(\bigcup \{Q_1[Y] \mid Q_1 \in \Delta_{1,\star}\} \right).$$

Clearly, $\mathcal{S}_1^\#$ is an extended saturated multiplicative system of $D_1[Y]$, and so we can consider the stable semistar operation of finite type on D_1 , $\circ_{\mathcal{S}_1^\#}$, defined by setting for each $A \in \overline{F}(D_1) = \overline{F}(D[X])$

$$A^{\circ_{\mathcal{S}_1^\#}} := AD_1[Y]_{\mathcal{S}_1^\#} \cap K_1.$$

We have already observed in Remark 19(b) that $[\tilde{\star}]$ is the unique stable semistar operation on $D_1 (= D[X])$ determined by the localizing system of finite type $\mathcal{F}_1 := \mathcal{F}_{1,\star} := \mathcal{F}^{\star_f}[X] = \mathcal{F}^\star[X]$. Moreover, the map $\sigma : \mathcal{F}_1 \mapsto \mathcal{S}(\mathcal{F}_1) := D_1[Y] \setminus \bigcup \{P_1[Y] \mid P_1 \notin \mathcal{F}_1\}$ establishes a bijection between the set of the localizing systems of finite type on D_1 and the extended saturated multiplicative systems of

$D_1[Y]$ and, under this map, the corresponding associated stable semistar operations of finite type on D_1 coincide [3, Corollary 2.2]. Since it is straightforward that $\Delta_{1,\star}$ coincides with the set $\{P_1 \in \text{Spec}(D_1) \mid P_1 \notin \mathcal{F}_{1,\star}\}$, then clearly $\mathcal{S}_{1,\star}^\#$ corresponds canonically to $\mathcal{F}_{1,\star}$ under σ . Therefore, $[\tilde{\star}]$ coincides with $\odot_{\mathcal{S}_{1,\star}^\#}$. Moreover, we also have $\text{Na}(D_1, [\tilde{\star}]) = D_1[Y]_{\mathcal{S}_{1,\star}^\#}$ or, equivalently, $\mathcal{S}_{1,\star}^\# = \{0 \neq g_1 \in D_1[Y] \mid c_{D_1}(g_1)^{[\tilde{\star}]} = D_1^{[\tilde{\star}]}\} =: \mathcal{N}_1^{[\tilde{\star}]}$ [3, Theorem 2.1(c) and (d)].

Let $\nabla_{1,\star}$ be the set of the maximal elements of $\Delta_{1,\star}$.

It is easy to see that

$$\begin{aligned} \nabla_{1,\star} = \{Q_1 \in \text{Spec}(D_1) \mid & \text{either } Q_1 \cap D = (0) \text{ and } c_D(Q_1)^{\star_f} = D^{\star_f} \\ & \text{or } Q_1 \cap D \in \text{QMax}^{\star_f}(D)\}, \end{aligned}$$

since a prime ideal $Q_1 \in \text{Spec}(D_1)$ such that $Q_1 \cap D = (0)$ is not contained in any ideal of the type $Q[X]$ with $Q \in \text{QMax}^{\star_f}(D)$ if and only if $c_D(Q_1)^{\star_f} = D^{\star_f}$.

For the sake of simplicity, set $\star_1 := \odot_{\mathcal{S}_{1,\star}^\#} (= [\tilde{\star}])$. Then, by the previous remarks, we can conclude that $\text{QMax}^{\star_1}(D_1) = \nabla_{1,\star}$ [3, Theorem 2.1(e)].

Putting together the previous information with Proposition 16(1), for all $A \in \overline{F}(D[X])$, we have

$$\begin{aligned} \bigcap \{AD_Q[X] \mid Q \in \text{QMax}^{\star_f}(D)\} &= A^{[\tilde{\star}]} = \bigcap \{AD[X]_{Q_1} \mid Q_1 \in \nabla_{1,\star}\} \\ &= AD[X, Y]_{\mathcal{N}_1^{[\tilde{\star}]}} \cap K(X) \\ &= A\text{Na}(D[X], [\tilde{\star}]) \cap K(X). \end{aligned}$$

In particular, for all $E \in \overline{F}(D)$,

$$\begin{aligned} E\text{Na}(D, \star) \cap K &= E^{\tilde{\star}} = E^{\tilde{\star}}[X] \cap K \\ &= (E[X])^{[\tilde{\star}]} \cap K \\ &= (E\text{Na}(D[X], [\tilde{\star}]) \cap K(X)) \cap K \\ &= E\text{Na}(D[X], [\tilde{\star}]) \cap K. \end{aligned}$$

- (2) Like $[\tilde{\star}]$, also $\langle \tilde{\star} \rangle$, and $(\tilde{\star})$ are finite type stable semistar operations on $D_1 := D[X]$ (Propositions 16(2) and 17) then, for the following multiplicative subsets of D_1 ,

$$\begin{aligned} \mathcal{S}_1^\#(\langle \tilde{\star} \rangle) &:= \mathcal{N}_1^{\langle \tilde{\star} \rangle} := \{0 \neq g_1 \in D_1[Y] \mid c_{D_1}(g_1)^{\langle \tilde{\star} \rangle} = D_1^{\langle \tilde{\star} \rangle}\}, \\ \mathcal{S}_1^\#((\tilde{\star})) &:= \mathcal{N}_1^{(\tilde{\star})} := \{0 \neq g_1 \in D_1[Y] \mid c_{D_1}(g_1)^{(\tilde{\star})} = D_1^{(\tilde{\star})}\}, \end{aligned}$$

and, for all $A \in \overline{F}(D[X])$, using also Proposition 16(1), we have

$$\begin{aligned} A^{\langle \tilde{\star} \rangle} &= AD[X, Y]_{\mathcal{N}_1^{\langle \tilde{\star} \rangle}} \cap K(X) = A\text{Na}(D[X], \langle \tilde{\star} \rangle) \cap K(X) \\ &= A\text{Na}(D, \star) \cap AK[X], \end{aligned}$$

$$\begin{aligned} A^{(\star)} &= AD[X, Y]_{\mathcal{N}_1^{(\star)}} \cap K(X) = A\text{Na}(D[X], (\tilde{\star})) \cap K(X) \\ &= A\text{Na}(D, \star). \end{aligned}$$

In particular, for all $E \in \overline{F}(D)$, we have

$$\begin{aligned} E\text{Na}(D, \star) \cap K &= E^{\tilde{\star}} = E^{\tilde{\star}}[X] \cap K = (E[X])^{(\tilde{\star})} \cap K = E\text{Na}(D[X], \langle \tilde{\star} \rangle) \cap K \\ &= (E[X])^{(\star)} \cap K = E\text{Na}(D[X], (\tilde{\star})) \cap K, \end{aligned}$$

with $(E[X])^{(\star)} = E\text{Na}(D, \star)$.

Note that, by the previous descriptions of $\langle \tilde{\star} \rangle$ and $(\tilde{\star})$, we have

$$\begin{aligned} P_1 \in \text{QSpec}^{(\star)}(D[X]) &\Leftrightarrow P_1\text{Na}(D, \star) \cap P_1K[X] \cap D[X] = P_1, \\ P_1 \in \text{QSpec}^{(\star)}(D[X]) &\Leftrightarrow P_1\text{Na}(D, \star) \cap D[X] = P_1. \end{aligned}$$

Since $\text{Na}(D, \star) = D[X]_{\mathcal{N}^{\star}}$, where $\mathcal{N}^{\star} := \{g \in D[X] \mid 0 \neq g \text{ and } c_D(g)^{\star} = D^{\star}\}$, then

$$\begin{aligned} \text{QSpec}^{(\star)}(D[X]) &= \{P_1 \in \text{Spec}(D[X]) \mid c_D(P_1)^{\star_f} \subsetneq D^{\star} \text{ or } P_1 \cap D = (0)\}, \\ \text{QSpec}^{(\star)}(D[X]) &= \{P_1 \in \text{Spec}(D[X]) \mid c_D(P_1)^{\star_f} \subsetneq D^{\star}\}. \end{aligned}$$

Therefore, since $\text{Max}(\text{Na}(D, \star)) = \{Q[X]_{\mathcal{N}^{\star}} \mid Q \in \text{QMax}^{\star_f}(D)\}$ [10, Proposition 3.1(2) and (3)], we can conclude that

$$\begin{aligned} \text{QMax}^{(\star)}(D[X]) &= \{Q_1 \in \text{Spec}(D[X]) \mid Q_1 = Q[X] \text{ for some } Q \in \text{QMax}^{\star_f}(D)\} \cup \\ &\quad \{(0) \neq Q_1 \in \text{Spec}(D[X]) \mid Q_1 \cap D = (0) \text{ and } c_D(Q_1)^{\star_f} = D^{\star}\}, \\ \text{QMax}^{(\star)}(D[X]) &= \{Q_1 \in \text{Spec}(D[X]) \mid Q_1 = Q[X] \text{ for some } Q \in \text{QMax}^{\star_f}(D)\}. \end{aligned}$$

We already observed that the construction described in Remark 20(1) is a modification of a previous construction due to Chang and Fontana [3, Theorem 2.3]. More precisely, Chang and Fontana considered the following subset of $\text{Spec}(D_1)$:

$$\begin{aligned} \Delta'_1 &:= \Delta'_{1,\star} := \{Q_1 \in \text{Spec}(D_1) \mid \text{either } Q_1 \cap D = (0) \text{ or} \\ &\quad Q_1 = (Q_1 \cap D)[X] \text{ and } (Q_1 \cap D)^{\star_f} \subsetneq D^{\star}\}. \end{aligned}$$

Then, they considered the following associated extended saturated multiplicative system in $D_1[Y]$:

$$\mathcal{S}'_1 := \mathcal{S}'_{1,\star} := \mathcal{S}(\Delta'_{1,\star}) := D_1[Y] \setminus \left(\bigcup \{Q_1[Y] \mid Q_1 \in \Delta'_{1,\star}\} \right)$$

and the stable semistar operation of finite type of D_1 defined by

$$A^{\circ_{\mathcal{S}'_{1,\star}}} := AD_1[Y]_{\mathcal{S}'_1} \cap K_1, \quad \text{for all } A \in \overline{F}(D_1).$$

They proved that, when \star is the v -operation (or the t -operation, or the w -operation) on D , then $\circ_{\mathcal{S}'_{1,\star}}$ coincides with the w -operation of $D[X]$ [3, Theorem 2.3(f)]. (Note

that, in that article, the authors denoted the semistar operation $\odot_{\mathcal{S}'_1, \star}$ of $D[X]$ by $[\star]$; we avoid now this notation, since we already use it here with a different meaning.)

Proposition 21. *Let \star be a semistar operation defined on an integral domain D . Let $\star'_1 := \odot_{\mathcal{S}'_1, \star}$ be the stable semistar operation of finite type on $D[X]$ defined above, and let \star_1 be the semistar operation defined in Remark 20(1) (i.e., $\star_1 := \odot_{\mathcal{S}'_1, \star} = [\star]$).*

- (1) $[\star] = \star_1 \leq \star'_1$.
- (2) $\star'_1 = \langle \star \rangle$.

Proof. (1) follows easily from the fact that $\Delta_{1, \star} \supseteq \Delta'_{1, \star}$ or, equivalently, $\mathcal{S}'_1 \subseteq \mathcal{S}_1$.

(2) We know that $\text{QMax}^{\star'_1}(D_1) = \{Q_1 \in \text{Spec}(D_1) \mid Q_1 \cap D = (0) \text{ and } c_D(Q_1)^{\star'} = D^*\} \cup \{Q[X] \mid Q \in \text{QMax}^{\star'}(D)\}$ [3, Theorem 2.3(e)]. On the other hand, by what we observed in Remark 20(2),

$$\text{QMax}^{\star'_1}(D[X]) = \text{QMax}^{\langle \star \rangle}(D[X]),$$

and so, since \star'_1 and $\langle \star \rangle$ are both stable semistar operations of finite type, we conclude that $\star'_1 = \langle \star \rangle$. \square

As a final remark (with the notation of the present article), note that in [3, Corollary 2.5(1)] the authors prove that D is a Prüfer \star -multiplication domain if and only if $D[X]$ is a Prüfer $\langle \star \rangle$ -multiplication domain. On the other hand, it is not true that D is a Prüfer \star -multiplication domain if and only if $D[X]$ is a Prüfer $[\star]$ -multiplication domain (take, for instance, D a Prüfer domain, but not a field, and $\star = d_D$, in this case $[\star] = d_{D[X]} = d_D$ and, obviously, $D[X]$ is not a Prüfer domain). This fact justifies the terminology used in [3], where the authors call the semistar operation denoted here by $\langle \star \rangle$ the stable semistar operation of finite type canonically associated to \star .

ACKNOWLEDGMENTS

We thank Muhammad Zafrullah for the helpful suggestions received during the preparation of the present article and the referee for providing constructive comments.

The second author was partially supported by a MIUR-PRIN grant no. 2008WYH9NY.

REFERENCES

- [1] Anderson, D. D. (1988). Star-operations induced by overrings. *Comm. Algebra* 16:2535–2553.
- [2] Bourbaki, N. (1961). *Algèbre Commutative*. Chapitres 1–2, Paris: Hermann.
- [3] Chang, G. W., Fontana, M. (2007). Uppers to zero and semistar operations in polynomial rings. *J. Algebra* 318:484–493.
- [4] Chang, G. W., Fontana, M. (2009). Uppers to zero in polynomial rings and Prüfer-like domains. *Comm. Algebra* 37:164–192.

- [5] El Baghdadi, S., Fontana, M. (2004). Semistar linkedness and flatness, Prüfer semistar multiplication domains. *Comm. Algebra* 32:1101–1126.
- [6] Fontana, M., Huckaba, J. A. (2000). Localizing systems and semistar operations. In: Chapman, S. T., Glaz, S., eds. *“Non-Noetherian Commutative Ring Theory”*. Kluwer Academic Publishers, pp. 169–198.
- [7] Fontana, M., Huckaba, J. A., Papick, I. J. (1997). *Prüfer Domains*. New York: Marcel Dekker Inc.
- [8] Fontana, M., Loper, K. A. (2001). A Krull-type theorem for the semistar integral closure of an integral domain. *AJSE (Arab J. Sci. Eng.) Theme Issue Commutative Algebra* 26:89–95.
- [9] Fontana, M., Loper, K. A. (2001). Kronecker function rings: a general approach. In: *Ideal Theoretic Methods in Commutative Algebra*. Lecture Notes in Pure Appl. Math. 220. Marcel Dekker, pp. 189–205.
- [10] Fontana, M., Loper, K. A. (2003). Nagata rings, Kronecker function rings and related semistar operations. *Comm. Algebra* 31:4775–4801.
- [11] Fontana, M., Loper, K. A. (2009). Cancellation properties in ideal systems: A classification of \mathbf{eab} semistar operations. *J. Pure Appl. Algebra* 213:2095–2103.
- [12] Gilmer, R. (1972). *Multiplicative Ideal Theory*. New York: Marcel Dekker.
- [13] Glaz, S., Vasconcelos, W. V. (1977). Flat ideals, II. *Manuscripta Math.* 22:325–341.
- [14] Halter-Koch, F. (1998). *Ideal Systems, An Introduction to Multiplicative Ideal Theory*. New York: Marcel Dekker.
- [15] Halter-Koch, F. (1997). Generalized integral closures. In: Anderson, D. D., ed. *Factorization in Integral Domains*. Vol. 187. Lecture Notes in Pure Appl. Math., Marcel Dekker, pp. 349–358.
- [16] Halter-Koch, F. (1998). *Ideal Systems, Monographs and Textbooks in Pure and Applied Mathematics*. Vol. 211. New York: Marcel Dekker Inc.
- [17] Halter-Koch, F. (2001). Localizing systems, module systems and semistar operations. *J. Algebra* 238:723–761.
- [18] Halter-Koch, F. (2003). Kronecker function rings and generalized integral closures. *Comm. Algebra* 31:45–59.
- [19] Hedstrom, J. R., Houston, E. (1980). Some remarks on star-operations. *J. Pure Appl. Algebra* 18:37–44.
- [20] Houston, E., Malik, S., Mott, J. (1984). Characterizations of $*$ -multiplication domains. *Canad. Math. Bull.* 27:48–52.
- [21] Houston, E., Zafrullah, M. (1989). On t -invertibility, II. *Comm. Algebra* 17:1955–1969.
- [22] Jaffard, P. (1960). *Les Systèmes d’Idéaux*. Paris: Dunod.
- [23] Kang, B. G. (1989). Prüfer v -multiplication domains and the ring $R[X]_{N_v}$. *J. Algebra* 123:151–170.
- [24] Lorenzen, P. (1939). Abstrakte Begründung der multiplicativen Idealtheorie. *Math. Z.* 45:533–553.
- [25] Matsumura, H. (1986). *Commutative Ring Theory*. Cambridge: Cambridge University Press.
- [26] Mimouni, A. (2008). Note on star operations over polynomial rings. *Comm. Algebra* 36:4249–4256.
- [27] Okabe, A., Matsuda, R. (1992). Star operations and generalized integral closures. *Bull. Fac. Sci. Ibaraki Univ. Ser. A* 24:7–13.
- [28] Okabe, A., Matsuda, R. (1994). Semistar operations on integral domains. *Math. J. Toyama Univ.* 17:1–21.
- [29] Picozza, G. (2004). *Semistar Operations and Multiplicative Ideal Theory*. Ph.D. thesis, Rome, Italy: Università degli Studi “Roma Tre”.
- [30] Picozza, G. (2007). A note on semistar Noetherian domains. *Houston J. Math.* 33:415–432.

- [31] Popescu, N. (1973). *Abelian categories with Applications to Rings and Modules*. L. M. S. Monograph No. 3. London: Academic Press.
- [32] Querré, J. (1980). Idéaux divisoriels d'un anneau de polynômes. *J. Algebra* 64:270–284.
- [33] Sahandi, P. (2009). Semistar-Krull and valuative dimension of integral domains. *Ricerche Mat.* 58:219–242.
- [34] Uda, H. (1983). LCM-stableness in ring extensions. *Hiroshima Math. J.* 13:357–377.
- [35] Wang, F. G., MacCasland, R. L. (1997). On w -modules over strong Mori domains. *Comm. Algebra* 25:1285–1306.
- [36] Zariski, O., Samuel, P. (1991). *Commutative Algebra*. Vol. II. New York: Van Nostrand.